

Black Vine Weevil

Introduction

The black vine weevil (*Otiorhynchus sulcatus*) may be found in herbaceous perennial production nurseries. It has a wide host range, feeding on over 200 plant species and may be spread by the transport of infected plant material.

Feeding Damage

Adults feed at night and hide around the base of plants during the day. As the adults feed, they cause c-shaped notching along the leaf margin to both herbaceous and woody plants. This damage is primarily cosmetic and may be hard to see on certain plants such as *Astilbe*.



Figure 1: Leaf notching at edges of leaves caused by adult black vine weevil. Photos by L. Pundt

The young larvae feed on fine roots and the mature larvae feed on stems, crowns, and larger roots. In herbaceous perennials, the larvae often feed in the crown severely reducing root systems and causing plant death.

Some of the perennials prone to damage include *Astilbe*, *Bergenia*, *Epimedium*, *Helleborus*, *Heuchera*, *Heucherella*, *Hosta*, *Physostegia*, *Phlox*, *Primula*, *Saxifrage*, *Sedum* and *Tricyrtis*. Some herbaceous perennials are very good hosts, such as *Astilbe* or *Tricyrtis*, but seem to tolerate more damage than other species.

Black vine weevils can also enter greenhouses and have been reported to damage some greenhouse crops such as begonia, impatiens, gloxinia, geranium and roses.




Figure 2: Mature larvae fed upon Heuchera crown and roots, causing wilting and plant death. Photos by L. Pundt

Biology and Life Cycle

Their life cycle progresses from egg, to larva, pupa, and adult. The black vine weevil adult is 3/8-inch-long, brown-to-black in color with yellow-brown markings on its wing covers, with ridges extending down the length of the abdomen. Adults have a distinct, short snout and elbowed antennae. Adult black vine weevils cannot fly because their wing covers are fused together.



Figure 3: Adult black vine weevil. Photo by R. Cowles (on left) and black vine weevil larvae (Photo by L. Pundt (on right)

After approximately two to three weeks of adult feeding, egg laying begins. Outdoors, adult female weevils lay eggs from July through August. Adults can live more than one year, often laying 200 to 400 eggs the first year and over 400 eggs in the second year. Eggs hatch in about two weeks and are laid in the upper one to two inches of the growing media. All adults are females who do not need to mate to reproduce.

Larvae hatch in August and begin feeding on fine young roots. Larvae are white to cream colored, legless grubs with yellow-brown heads and are covered with fine hairs. The larvae have six instars. When mature, they are about 1/2 inch long. Older larvae are also slightly C-shaped but lack legs. Black vine weevil larvae feed on the roots of host plants and then overwinter as full-grown larvae. Larvae resume feeding in the

early spring, often causing the heaviest damage. Pupation occurs in earthen cells that they form in the growing medium.

There is generally one generation per year, but in greenhouses, depending upon temperatures and environmental conditions, several overlapping generations with different life stages may be present.

Scouting

Since the larvae are active in the growing media and adults feed at night by notching the edges of leaves, infestations may go unnoticed until significant damage has occurred. Although adults are active at night, they may be monitored by placing horizontal sticky boards on benches to trap adults. Check plant debris and growing medium around plants for the presence of adult black vine weevils. During the day, adults stay at the base of favored host plants, under and on the side of containers. Adults may be active during cloudy, overcast days.

Larvae feed on roots, causing wilting as roots are damaged. If plants exhibit these symptoms, check the root system and surrounding root ball for the presence of larvae.

Biological Controls

Black vine weevil larvae are susceptible to nearly all commercially available species of entomopathogenic nematodes. Some of the commercially available species include *Heterorhabditis bacteriophora*, *Steinernema kraussei* (Nemasys L) and *Steinernema carpocapsae* that are applied as drenches to the growing medium. Apply the nematodes to moist soil in the early morning or evening to avoid heat and direct sunlight. The nematodes are attracted to, enter, and kill all larval stages and pupae, but not eggs or adults.

Heterorhabditis bacteriophora is applied to the soil with a minimum temperature of 57° F. The infected larvae turn brick red to maroon. *Steinernema kraussei* is effective in a wide range of temperatures from 41 to 86° F; infected larvae or pupae turn dark yellow. Consider other *Steinernema* species if soil temperatures exceed 60°F. Infected larvae or pupae remain creamy yellow-white, but you can see nematode movement through the cuticle with a hand lens. These nematodes can be applied from March to May and from August until November to target the larvae after eggs have been laid. The nematodes are active up to 6 weeks after their application. Repeated applications may be needed depending upon the level of infestation. Nematodes should be applied to moist growing media during cloudy, overcast weather.

Cultural Controls

Adults cannot fly but may invade heated structures to overwinter or may enter outdoor production areas from surrounding vegetation or be accidentally brought in on infested plant material.

- Avoid black vine weevil infestations by buying nursery stock certified as weevil-free.
- Closely inspect incoming plants to exclude weevils from production areas.
- Segregate infested plant material from susceptible new plantings.

- Check older stock material when dividing.
- Handpick adults if only a few plants are infested.
- Immediately discard heavily infested plants.
- Hedgerows of yew, arborvitae or spruce surrounding production areas are good hosts of black vine weevils and may be a source of an infestation.

Chemical Controls

It may be necessary to apply an insecticide drench to protect highly susceptible plants. Drench applications of the insecticide, bifenthrin (Talstar N Granular®) effectively kills black vine weevil larvae in containers and in the growing medium. It is applied as a pre-plant potting mix incorporation based upon the known bulk density of the media as a preventative treatment. See label for special use instructions. All the soil volume must be treated.

By: Leanne Pundt, Extension Educator, UConn Extension, 2016, latest revision 2024

References

- Cloyd, R. 2015. Weeding out the Weevil. Greenhouse Management. <http://www.greenhousemag.com/article/gm0115-active-black-vine-weevil-control/>
- Cowles, R. 2003. Practical Black Vine Weevil Management. Journal of American Rhododendron Society (JARS). 57(4): 219.
- Cowles, R. 2001. Protecting Container-Grown Crops from Black Vine Weevil Larvae with Bifenthrin. J. Environ. Hort. 19(4):184-189.
- Gill, S., J. Lutz, P. Shrewsbury, and M. Raupp. 2001. Evaluation of Biological and Chemical Control Methods for Black Vine Weevil, *Otiorhynchus sulcatus* (Fabricius) (Coleoptera: Curculionidae) in Container Grown Perennials. J. Environ. Hort. 19(3):166-170.
- Gill, S., and J. Sanderson. 1998. Ball Identification Guide to Greenhouse Pests and Beneficials. Ball Publishing. Batavia, Ill. 244 pp.
- Gill, S. and P. Shrewsbury. 2013. Black Vine Weevil: Controlling a Major Nursery and Landscape Pest. University of Maryland FS – 805. Updated 2021.
- Georgis, R. and G. Poinar. 1984. Greenhouse Control of the Black Vine Weevil *Otiorhynchus sulcatus* (Coleoptera: Curculionidae) by Heterorhabditid and Steinernematid Nematodes. Environ. Entomol. 13: 1138-1140.

Disclaimer for Fact Sheets:

The information in this document is for educational purposes only. The recommendations contained are based on the best available knowledge at the time of publication. Any reference to commercial products, trade or brand names is for information only, and no endorsement or approval is intended. UConn Extension does not guarantee or warrant the standard of any product referenced or imply approval of the product to the exclusion of others which also may be available. The University of Connecticut, UConn Extension, College of Agriculture, Health and Natural Resources is an equal opportunity program provider and employer.